

Author Index

Abe, K., Kawagoe, J., Lee, T.-H., Aoki, M. and Kogure, K.
Disturbance of a mitochondrial DNA expression in gerbil hippocampus after transient forebrain ischemia (19) 69

Aceves, J., see Arias-Montano, J.-A. (19) 233

Agarwal, N., see Nir, I. (19) 47

Akagawa, K., see Inoue, A. (19) 121

Alkon, D.L., see Craig, A.M. (19) 269

Aoki, M., see Abe, K. (19) 69

Arias-Montano, J.-A., Aceves, J. and Young, J.M.
Carbachol-induced phosphoinositide metabolism in slices of rat substantia nigra pars reticulata (19) 233

Asanuma, M., see Chou, H.-h. (19) 211

Azmitia, E.C., see Liao, B. (19) 328

Baldessari, S., see Bendotti, C. (19) 257

Barizo, M.J., see Sanna, P.P. (19) 241

Baudry, M., see Wei, W. (19) 339

Bendotti, C., Baldessari, S., Ehret, M., Tarizzo, G. and Samanin, R.
Effect of d-fenfluramine and 5,7-dihydroxytryptamine on the levels of tryptophan hydroxylase and its mRNA in rat brain (19) 257

Bengzon, J., see Kokaia, Z. (19) 277

Bloom, F.E., see Sanna, P.P. (19) 241

Bogic, L., see Hammer Jr., R.P. (19) 129

Bourbonnière, M. and Nalbantoglu, J.
Expression of amyloid precursor protein in a neuronal cell line: functional activity of proximal regulatory elements (19) 246

Bredesen, D.E., see Zhong, L.-t. (19) 353

Broek, D., see Wei, W. (19) 339

Brooks, P.J., Funabashi, T., Kleopoulos, S.P., Mobbs, C.V. and Pfaff, D.W.
Cell-specific expression of preproenkephalin intronic heteronuclear RNA in the rat forebrain (19) 22

Burgess, L.H. and Handa, R.J.
Hormonal regulation of androgen receptor mRNA in the brain and anterior pituitary gland of the male rat (19) 31

Byrne, J.H., see Noel, F. (19) 203

Chou, H.-h., Ogawa, N., Asanuma, M., Hirata, H., Kondo, Y. and Mori, A.
Rapid response of striatal muscarinic M₁-receptor mRNA to muscarinic cholinergic agents in rat brain (19) 211

Chritin, M., Feuerstein, C. and Savasta, M.
Time-course of changes in striatal levels of DA uptake sites, DA D₂ receptor and preproenkephalin mRNAs after nigrostriatal dopaminergic denervation in the rat (19) 318

Chua Jr., S.C., see Jhanwar-Uniyal, M. (19) 195

Cook, R., see Noel, F. (19) 203

Craig, A.M., Olds, J.L., Schreurs, B.G., Scharenberg, A.M. and Alkon, D.L.
Quantitative distribution of protein kinase C α , β , γ , and ϵ mRNAs in the hippocampus of control and nictitating membrane conditioned rabbits (19) 269

De Blas, A., see Luntz-Leybman, V. (19) 9

De Groen, P.C., see Nielander, H.B. (19) 293

Devi, L., see Grigoriants, O. (19) 161

Donnelly-Roberts, D.L. and Lentz, T.L.
Sodium dodecyl sulfate- and carbamyl-choline-induced changes in circular dichroism spectra of acetylcholine receptor synthetic peptides (19) 55

Dreyer, D., see Pennypacker, K.R. (19) 349

Duman, R.S., see Vaccarino, F.M. (19) 76

Eggen, B.J.L., see Nielander, H.B. (19) 293

Ehret, M., see Bendotti, C. (19) 257

Elizarov, S.M. and Preobrazhensky, A.A.
Phosphorylation of two human neurochordins by mammalian casein kinase 1 (19) 310

Elliott, K., see Franklin, S.O. (19) 93

Eskin, A., see Noel, F. (19) 203

Fei, J., see Lam, D.M.-K. (19) 227

Feng, P., see Satoh, T. (19) 175

Feng, X., see Katagiri, T. (19) 1

Fernandez, J., see Schuligoi, R. (19) 333

Fernando, L., see Luntz-Leybman, V. (19) 9

Feuerstein, C., see Chritin, M. (19) 318

Fischer-Colbrie, R., see Mahata, S.K. (19) 83

Folsom, D.P., see Sanna, P.P. (19) 241

Franklin, S.O., Elliott, K., Zhu, Y.-S., Wahlestedt, C. and Inturrisi, C.E.
Quantitation of NMDA receptor (NMDAR1) mRNA levels in the adult and developing rat CNS (19) 93

Fricker, L.D., see Grigoriants, O. (19) 161

Frostholm, A., see Luntz-Leybman, V. (19) 9

Fujita, M., Shimada, S., Nishimura, T., Uhl, G.R. and Tohyama, M.
Ontogeny of dopamine transporter mRNA expression in the rat brain (19) 222

Funabashi, T., see Brooks, P.J. (19) 22

Furukawa, T., see Shibata, K. (19) 115

Furuyama, T., Inagaki, S. and Takagi, H.
Distribution of type II adenylyl cyclase mRNA in the rat brain (19) 165

Gidö, G., see Kokaia, Z. (19) 277

Gispert, W.H., see Nielander, H.B. (19) 293

Gray, C.W. and Patel, A.J.
Regulation of β -amyloid precursor protein isoform mRNAs by transforming growth factor- β 1 and interleukin-1 β in astrocytes (19) 251

Grigoriants, O., Devi, L. and Fricker, L.D.
Dopamine antagonist haloperidol increases carboxypeptidase E mRNA in rat neurointermediate pituitary but not in various other rat tissues (19) 161

Guo, L.-H., see Lam, D.M.-K. (19) 227

Hammer Jr., R.P., Bogic, L. and Handa, R.J.
Estrogenic regulation of proenkephalin mRNA expression in the ventromedial hypothalamus of the adult male rat (19) 129

Handa, R.J., see Burgess, L.H. (19) 31

Handa, R.J., see Hammer Jr., R.P. (19) 129

Hartigan, D.J., see Vaccarino, F.M. (19) 76

Hayward, M.D., see Vaccarino, F.M. (19) 76

Heavens, R.P., see Schuligoi, R. (19) 333

Hirata, H., see Chou, H.-h. (19) 211

Hirsch, M.D., see Sanna, P.P. (19) 241

Hockfield, S., see Martin, K.A. (19) 303

Hong, J.S., see Pennypacker, K.R. (19) 349

Horn, D.A. and Latchman, D.S.
The tissue specific SmN protein does not influence the alternative splicing of endogenous N-Cam and C-SRC RNAs in transfected 3T3 cells (19) 181

Huang, F., see Lam, D.M.-K. (19) 227

Iadarola, M.J., see Jian, M. (19) 156

Ichikawa, T., see Katagiri, T. (19) 1

Ihara, Y., see Kobayashi, H. (19) 188

Inagaki, S., see Furuyama, T. (19) 165

Inoue, A. and Akagawa, K.
Neuron specific expression of a membrane protein, HPC-1: tissue distribution, and cellular and subcellular localization of immunoreactivity and mRNA (19) 121

Inturrisi, C.E., see Franklin, S.O. (19) 93

Jhanwar-Uniyal, M. and Chua Jr., S.C.
Critical effects of aging and nutritional state on hypothalamic neuropeptide Y and galanin gene expression in lean and genetically obese Zucker rats (19) 195

Jian, M., Staines, W.A., Iadarola, M.J. and Robertson, G.S.
Destruction of the nigrostriatal pathway increases Fos-like immunoreactivity predominantly in striatopallidal neurons (19) 156

Johnson, G., Refolo, L.M., Merrill, C.R. and Wallace, W.
Altered expression and phosphorylation of amyloid precursor protein in heat shocked neuronal PC12 cells (19) 140

Johnson, G., see Wallace, W. (19) 149

Kalivas, P.W., see See, R.E. (19) 219

Kane, D.J., see Zhong, L.-t. (19) 353

Katagiri, T., Feng, X., Ichikawa, T., Usui, H., Takahashi, Y. and Kumanishi, T.
Neuron-specific enolase (NSE) and non-neuronal enolase (NNE) mRNAs are co-

expressed in neurons of the rat cerebellum: *in situ* hybridization histochemistry (19) 1

Kawagoe, J., see Abe, K. (19) 69

Kawasaki, M., Yoshihara, Y., Yamaji, M. and Watanabe, Y. Expression of prostaglandin endoperoxide synthase in rat brain (19) 39

Kim, U.J., see Satoh, T. (19) 175

King, S.C., see Lam, D.M.-K. (19) 227

Kiyama, H., Sato, K. and Tohyama, M. Characteristic localization of non-NMDA type glutamate receptor subunits in the rat pituitary gland (19) 262

Kiyama, H., Sato, K., Kuba, T. and Tohyama, M. Sympathetic and parasympathetic ganglia express non-NMDA type glutamate receptors: distinct receptor subunit composition in the principle and SIF cells (19) 345

Kiyama, H., see Saika, T. (19) 237

Kleopoulos, S.P., see Brooks, P.J. (19) 22

Kobayashi, H., Uchida, Y., Ihara, Y., Nakajima, K., Kohsaka, S., Miyatake, T. and Tsuji, S. Molecular cloning of rat growth inhibitory factor cDNA and the expression in the central nervous system (19) 188

Kogure, K., see Abe, K. (19) 69

Kohsaka, S., see Kobayashi, H. (19) 188

Kojima, N., Shirao, T. and Obata, K. Molecular cloning of a developmentally regulated brain protein, chicken drebrin A and its expression by alternative splicing of the drebrin gene (19) 101

Kokaia, M., see Kokaia, Z. (19) 277

Kokaia, Z., Gidö, G., Ringstedt, T., Bengzon, J., Kokaia, M., Siesjö, B.K., Persson, H. and Lindvall, O. Rapid increase of BDNF mRNA levels in cortical neurons following spreading depression: regulation by glutamatergic mechanisms independent of seizure activity (19) 277

Komatsu, C., see Shibata, K. (19) 115

Kondo, H., see Sakagami, H. (19) 215

Kondo, Y., see Chou, H.-h. (19) 211

Kreek, M.J., see Spangler, R. (19) 323

Krukoff, T.L. and Zheng, Y. Effects of renal denervation and reinnervation on ganglionic gene expression of neurotransmitter proteins and *c-fos* in rat (19) 287

Kuba, T., see Kiyama, H. (19) 345

Kubo, T., see Saika, T. (19) 237

Kumanishi, T., see Katagiri, T. (19) 1

Lam, D.M.-K., Fei, J., Zhang, X.-Y., Tam, A.C.W., Zhu, L.-H., Huang, F., King, S.C. and Guo, L.-H. Molecular cloning and structure of the human (GABATHG) GABA transporter gene (19) 227

Latchman, D.S., see Horn, D.A. (19) 181

Le, H.N., see Vaccarino, F.M. (19) 76

Lee, T.-H., see Abe, K. (19) 69

Lentz, T.L., see Donnelly-Roberts, D.L. (19) 55

Liao, B., Miesak, B. and Azmitia, E.C. Loss of 5-HT_{1A} receptor mRNA in the dentate gyrus of the long-term adrenalectomized rats and rapid reversal by dexamethasone (19) 328

Lindvall, O., see Kokaia, Z. (19) 277

Luntz-Leybman, V., Frostholm, A., Fernando, L., De Blas, A. and Rotter, A. GABA_A/benzodiazepine receptor γ_2 subunit gene expression in developing normal and mutant mouse cerebellum (19) 9

Maciejewski-Lenoir, D., see Sanna, P.P. (19) 241

Maggi, A., see Pollio, G. (19) 135

Mahata, M., see Mahata, S.K. (19) 83

Mahata, S.K., Mahata, M., Fischer-Colbrie, R. and Winkler, H. Reserpine causes differential changes in the mRNA levels of chromogranin B, secretogranin II, carboxypeptidase H, alpha-amidating monooxygenase, the vesicular amine transporter and of synaptophysin in rat brain (19) 83

Martin, K.A. and Hockfield, S. Expression of the *mas* proto-oncogene in the rat hippocampal formation is regulated by neuronal activity (19) 303

Matsunaga, T., see Saika, T. (19) 237

McMillian, M.K., see Pennypacker, K.R. (19) 349

Melia, K.R., see Sanna, P.P. (19) 241

Merril, C.R., see Johnson, G. (19) 140

Merril, C.R., see Wallace, W. (19) 149

Miesak, B., see Liao, B. (19) 328

Misumi, Y., see Shibata, K. (19) 115

Miyatake, T., see Kobayashi, H. (19) 188

Miyatake, T., see Saito, F. (19) 171

Mobbs, C.V., see Brooks, P.J. (19) 22

Morgan, J.I., see Sangameswaran, L. (19) 62

Mori, A., see Chou, H.-h. (19) 211

Nakajima, K., see Kobayashi, H. (19) 188

Nalbantoglu, J., see Bourbonnière, M. (19) 246

Nestler, E.J., see Vaccarino, F.M. (19) 76

Nicolin, A., see Pollio, G. (19) 135

Nielander, H.B., De Groen, P.C., Eggen, B.J.L., Schrama, L.H., Gispen, W.H. and Schotman, P. Structure of the human gene for the neural phosphoprotein B-50 (GAP-43) (19) 293

Nir, I. and Agarwal, N. Diurnal expression of *c-fos* in the mouse retina (19) 47

Nishimura, T., see Fujita, M. (19) 222

Noel, F., Nuñez-Regueiro, M., Cook, R., Byrne, J.H. and Eskin, A. Long-term changes in synthesis of intermediate filament protein, actin and other proteins in pleural sensory neurons of *Aplysia* produced by an in vitro analogue of sensitization training (19) 203

Nuñez-Regueiro, M., see Noel, F. (19) 203

Obata, K., see Kojima, N. (19) 101

Ogawa, N., see Chou, H.-h. (19) 211

Olds, J.L., see Craig, A.M. (19) 269

Patel, A.J., see Gray, C.W. (19) 251

Pennypacker, K.R., Dreyer, D., Hong, J.S. and McMillian, M.K. Elevated basal AP-1 DNA binding activity in developing rat brain (19) 349

Persson, H., see Kokaia, Z. (19) 277

Pfaff, D.W., see Brooks, P.J. (19) 22

Pollio, G., Xue, P., Zanisi, M., Nicolin, A. and Maggi, A. Antisense oligonucleotide blocks progesterone-induced lordosis behavior in ovariectomized rats (19) 135

Preobrazhensky, A.A., see Elizarov, S.M. (19) 310

Refolo, L.M., see Johnson, G. (19) 140

Refolo, L.M., see Wallace, W. (19) 149

Ringstedt, T., see Kokaia, Z. (19) 277

Robertson, G.S., see Jian, M. (19) 156

Rotter, A., see Luntz-Leybman, V. (19) 9

Saika, T., Takeda, N., Kiyama, H., Kubo, T., Tohyama, M. and Matsunaga, T. Changes in preproenkephalin mRNA after unilateral and bilateral labyrinthectomy in the rat medial vestibular nucleus (19) 237

Saito, F., Yanagisawa, K. and Miyatake, T. Soluble derivatives of β -A4 amyloid protein precursor in human cerebrospinal fluid are both *N*- and *O*-glycosylated (19) 171

Sakagami, H. and Kondo, H. Cloning and sequencing of a gene encoding the β polypeptide of Ca^{2+} /calmodulin-dependent protein kinase IV and its expression confined to the mature cerebellar granule cells (19) 215

Samanin, R., see Bendotti, C. (19) 257

Sangameswaran, L. and Morgan, J.I. Structure and regulation of the gene encoding the neuron-specific protein PEP-19 (19) 62

Sanna, P.P., Folsom, D.P., Barizo, M.J., Hirsch, M.D., Melia, K.R., Maciejewski-Lenoir, D. and Bloom, F.E. Chronic ethanol intake decreases vasopressin mRNA content in the rat hypothalamus: a PCR study (19) 241

Sato, K., see Kiyama, H. (19) 262

Sato, K., see Kiyama, H. (19) 345

Satoh, T., Feng, P., Kim, U.J. and Wilber, J.F. Identification of thyrotropin-releasing hormone receptor messenger RNA in the rat central nervous system and eye (19) 175

Savasta, M., see Chritin, M. (19) 318

Scharenberg, A.M., see Craig, A.M. (19) 269

Schotman, P., see Nielander, H.B. (19) 293

Schrama, L.H., see Nielander, H.B. (19) 293

Schreiber, S.S., see Wei, W. (19) 339

Schreurs, B.G., see Craig, A.M. (19) 269

Schuligoi, R., Fernandez, J., Heavens, R.P. and Sirinathsinghji, D.J.S.

Decreased tyrosine hydroxylase mRNA but not cholecystokinin mRNA in the pars compacta of the substantia nigra and ventral tegmental area of aged rats (19) 333

Sealfon, S.C., see Spyrosi, C. (19) 313

See, R.E., Striplin, C. and Kalivas, P.W. Chronic haloperidol does not alter G protein α -subunit levels in rats (19) 219

Shibata, K., Komatsu, C., Misumi, Y. and Furukawa, T. Developmental differences of angiotensinogen mRNA in the preoptic area between spontaneously hypertensive and age-matched Wistar-Kyoto rats (19) 115

Shimada, S., see Fujita, M. (19) 222

Shirao, T., see Kojima, N. (19) 101

Siesjö, B.K., see Kokaia, Z. (19) 277

Sirinathsinghji, D.J.S., see Schulgoi, R. (19) 333

Spangler, R., Unterwald, E.M. and Kreek, M.J. 'Binge' cocaine administration induces a sustained increase of prodynorphin mRNA in rat caudate-putamen (19) 323

Spyrosi, C. and Sealfon, S.C. Regulation of dopamine D₂ receptor mRNA expression in the olfactory tubercle by cocaine (19) 313

Staines, W.A., see Jian, M. (19) 156

Striplin, C., see See, R.E. (19) 219

Sugar, J., see Wallace, W. (19) 149

Takagi, H., see Furuyama, T. (19) 165

Takahashi, Y., see Katagiri, T. (19) 1

Takeda, N., see Saika, T. (19) 237

Tam, A.C.W., see Lam, D.M.-K. (19) 227

Tarizzo, G., see Bendotti, C. (19) 257

Tocco, G., see Wei, W. (19) 339

Tohyama, M., see Fujita, M. (19) 222

Tohyama, M., see Kiyama, H. (19) 262

Tohyama, M., see Kiyama, H. (19) 345

Tohyama, M., see Saika, T. (19) 237

Tsuji, S., see Kobayashi, H. (19) 188

Uchida, Y., see Kobayashi, H. (19) 188

Uhl, G.R., see Fujita, M. (19) 222

Unterwald, E.M., see Spangler, R. (19) 323

Usui, H., see Katagiri, T. (19) 1

Vaccarino, F.M., Hayward, M.D., Le, H.N., Hartigan, D.J., Duman, R.S. and Nestler, E.J. Induction of immediate early genes by cyclic AMP in primary cultures of neurons from rat cerebral cortex (19) 76

Wahlestedt, C., see Franklin, S.O. (19) 93

Wallace, W., Johnson, G., Sugar, J., Merrill, C.R. and Refolo, L.M.

Reversible phosphorylation of tau to form A68 in heat-shocked neuronal PC12 cells (19) 149

Wallace, W., see Johnson, G. (19) 140

Watanabe, Y., see Kawasaki, M. (19) 39

Wei, W., Schreiber, S.S., Baudry, M., Tocco, G. and Broek, D. Localization of the cellular expression pattern of *cdc25^{NEF}* and *ras* in the juvenile rat brain (19) 339

Wilber, J.F., see Satoh, T. (19) 175

Winkler, H., see Mahata, S.K. (19) 83

Xue, P., see Pollio, G. (19) 135

Yamaji, M., see Kawasaki, M. (19) 39

Yanagisawa, K., see Saito, F. (19) 171

Yoshihara, Y., see Kawasaki, M. (19) 39

Young, J.M., see Arias-Montano, J.-A. (19) 233

Zanisi, M., see Pollio, G. (19) 135

Zhang, X.-Y., see Lam, D.M.-K. (19) 227

Zheng, Y., see Krukoff, T.L. (19) 287

Zhong, L.-t., Kane, D.J. and Bredesen, D.E. BCL-2 blocks glutamate toxicity in neural cell lines (19) 353

Zhu, L.-H., see Lam, D.M.-K. (19) 227

Zhu, Y.-S., see Franklin, S.O. (19) 93

